

Finite-Temperature Mott Transition in the Two-Dimensional Hubbard Model

S. Onoda^{*a} and M. Imada^b

^aTokura Spin Superstructure Project, ERATO, Japan Science and Technology Corporation c/o, Department of Applied Physics, University of Tokyo, Tokyo 113-8656, Japan

^bInstitute for Solid State Physics, University of Tokyo, Kashiwa, Chiba 277-8581, Japan

Mott transitions are studied in the two-dimensional Hubbard model by a non-perturbative theory of correlator projection that systematically includes spatial correlations into the dynamical mean-field theory. A nonzero second-neighbor transfer yields a first-order Mott transition at finite temperatures ending at a critical point.

Keywords: Mott transition; Hubbard model

The Mott transition has been one of the fundamental issues in strongly correlated electrons [1]. In spite of substantial progress [1,2], its full understanding in low dimensions remains open.

Here, we propose a non-perturbative theoretical framework of correlator projection method (CPM) [3]. The CPM combines the operator projection method [4] with the dynamical mean-field approximation (DMFA) by restoring spatial correlations along the coarse-graining concept in the energy-momentum space with increasing the order of projection. It does not suffer from a cluster-size problem. Then, we obtain the Mott transition phase diagram of the two-dimensional (2D) Hubbard model in the parameter space of the Coulomb repulsion U , the second-neighbor transfer t' , the temperature T (scaled by the nearest-neighbor transfer t) and the filling $\langle n \rangle$. A first-order Mott transition surface with a critical end curve appears at half filling. Doping into the Mott insulator yields a diverging compressibility and/or a tendency to phase separation.

We consider the 2D Hubbard model. The formalism of CPM [3] is as follows: The operator projection leads to series of Dyson equations,

$$\Sigma_{n-1}(\omega, \mathbf{k}) = G_{n-1}(\omega, \mathbf{k}) \varepsilon_{\mathbf{k}}^{(n, n-1)} \quad (1)$$

^{*}Corresponding author. Tel.: +81-3-5841-7904; fax.: +81-3-5841-7904. E-mail address: sonoda@appi.t.u-tokyo.ac.jp

$$G_{n-1}(\omega, \mathbf{k}) = 1/[\omega - \varepsilon_{\mathbf{k}}^{(n, n)} - \Sigma_n(\omega, \mathbf{k})] \quad (2)$$

with $n = 1, 2, \dots$. $G_0 = \Sigma_0$ is the electron Green's function and Σ_1 is the self-energy part with $\varepsilon_{\mathbf{k}}^{(1,0)} = 1$. Other coefficients up to the second order $n = 2$ are given by the Hartree-Fock dispersion $\varepsilon_{\mathbf{k}}^{(1,1)} = -t_{\mathbf{k}} - \mu - U\langle n \rangle/2$, a source of Hubbard band splitting $\varepsilon_{\mathbf{k}}^{(2,1)} = U^2 M$ with $M = \langle n \rangle(2 - \langle n \rangle)/4$, and $\varepsilon_{\mathbf{k}}^{(2,2)} = -\tilde{t}_{\mathbf{k}} - \mu - U(1 - \langle n \rangle/2) - \varepsilon_{\text{cor}}/M$. Here, μ is the chemical potential determined from the filling $\langle n \rangle$. $\varepsilon_{\text{cor}} = -\sum_{\mathbf{x}', s} t_{\mathbf{x}, \mathbf{x}'} \langle (1/2 - n_{\mathbf{x}, -s})(c_{\mathbf{x}, s}^\dagger c_{\mathbf{x}', s} + h.c.) \rangle / 2$ is a local energy shift due to a correlated hopping process with $n_{\mathbf{x}, s} = c_{\mathbf{x}, s}^\dagger c_{\mathbf{x}, s}$ where $c_{\mathbf{x}, s}$ and $c_{\mathbf{x}, s}^\dagger$ are the electron creation and annihilation operators at a site $\mathbf{x}$ with a spin s , respectively. $t_{\mathbf{k}}$ and $\tilde{t}_{\mathbf{k}}$ are the Fourier transforms of the transfer $t_{\mathbf{x}, \mathbf{x}'}$ and $\tilde{t}_{\mathbf{x}, \mathbf{x}'} = t_{\mathbf{x}, \mathbf{x}'} (\langle n_{\mathbf{x}} n_{\mathbf{x}'} \rangle / 4 + \langle \tilde{S}_{\mathbf{x}} \cdot \tilde{S}_{\mathbf{x}'} \rangle - \langle \Delta_{\mathbf{x}}^\dagger \Delta_{\mathbf{x}'} \rangle) / M$, respectively, where $n_{\mathbf{x}}$, $\tilde{S}_{\mathbf{x}}$ and $\Delta_{\mathbf{x}}$ represent the charge, spin and local-pair operators, respectively. $\tilde{t}_{\mathbf{k}}$ introduces $\mathbf{k}$ dependences of $\Sigma_1(\omega, \mathbf{k})$ mainly through the superexchange process and produces antiferromagnetic precursors. Here, we determine it from Ref. [5].

To reproduce a Mott insulator in particle-hole asymmetric cases, the local dynamics of $\Sigma_2(\omega, \mathbf{k})$ must be obtained in a sufficiently correct manner. Then, instead of decoupling approximations [4,6], we adopt the following gen-

eralized DMFA: (i) We calculate a local *normalized* self-energy part $G_{1,\text{loc}}(\omega) = \frac{1}{N} \sum_{\mathbf{k}} G_1(\omega, \mathbf{k})$ from an arbitrary $\Sigma_{2,\text{loc}}(\omega)$ by replacing $\Sigma_2(\omega, \mathbf{k})$ with $\Sigma_{2,\text{loc}}(\omega)$ in Eq. (2), (ii) calculate the *Weiss self-energy part* $\mathcal{S}_1(\omega) = \varepsilon^{(2,1)} / [G_{1,\text{loc}}^{-1}(\omega) + \Sigma_{2,\text{loc}}(\omega)]$, and (iii) obtain the *Weiss Green's function* $\mathcal{G}_0(\omega, \mathbf{k}) = 1 / [\omega - \varepsilon_{\mathbf{k}}^{(1,1)} - \mathcal{S}_1(\omega)]$. (iv) Within the iterative perturbation scheme, we calculate a new $\Sigma_{2,\text{loc}}(\omega)$ as a Fourier transform of $\Sigma_{2,\text{loc}}(\tau) = \frac{1}{N^3} \sum_{\mathbf{k}, \mathbf{k}', \mathbf{q}} \Gamma_{\mathbf{k}, \mathbf{k}', \mathbf{q}} \mathcal{G}_0(\tau, \mathbf{k} + \mathbf{q}) \mathcal{G}_0(\tau, \mathbf{k}' - \mathbf{q}) \mathcal{G}_0(-\tau, \mathbf{k}')$ where $\Gamma_{\mathbf{k}, \mathbf{k}', \mathbf{q}} = 4[t_{\mathbf{k}'} - t_{\mathbf{k}' - \mathbf{q}} - t_{\mathbf{k} + \mathbf{q}} + t_{\mathbf{k}} + 4\varepsilon_{\text{cor}} / (\langle n \rangle (2 - \langle n \rangle))]^2 / (\langle n \rangle (2 - \langle n \rangle))$. This loop continues to (i) until convergence is reached.

This second-order CPM reproduces a Mott insulator at low T 's for $t' = 0$ at half filling and its single-particle spectra and momentum distributions show a remarkable similarity to numerical results [7]. As we introduce $t' \neq 0$ with U and $\langle n \rangle = 1$ being fixed, the system undergoes a first-order Mott metal-insulator transition (MIT) at a low T accompanied by a discontinuity in the double occupancy $\langle n_{\uparrow} n_{\downarrow} \rangle$. In the space of (U, t', T) , this first-order MIT boundary forms a surface [3]. Within the surface with U being fixed, the jump in $\langle n_{\uparrow} n_{\downarrow} \rangle$ decreases with increasing T and disappears above a critical temperature T_{cr} ; the first-order surface ends at a critical curve at $T > 0$. Then, without any spontaneous symmetry breaking as in two dimensions at $T > 0$, the Mott transition shows a similarity to a liquid-gas phase transition, as in Refs. [8,2]. The MIT boundary at $T \rightarrow 0$ agrees with numerical results [9].

Doping carriers into Mott insulators yields a metallic state characterized by an enhanced charge compressibility $\kappa = \partial \langle n \rangle / \partial \mu$. The present results of $\mu - \delta$ curves near half filling are shown in Fig. 1 with $\delta = 1 - \langle n \rangle$. The slope $\partial \mu / \partial \langle n \rangle$ represents $1/\kappa$. Then, from high T 's, κ increases with decreasing T and shows a divergence towards the filling-controll MIT at low T 's. Even an instability to a phase separation appears as emergence of $\kappa < 0$ at lower T 's. However, the issue if the phase separation is an artifact of the present approximation requires a further intensive study.

In summary, using the CPM that systematically includes spatial correlations into the DMFA, the MIT phase diagram of the 2D Hubbard model

has been clarified in the $(U, |t'|, T)$ space with a first-order MIT surface with a $T > 0$ critical end curve for $t' \neq 0$ at half filling as well as a diverging charge compressibility and/or a tendency to a phase separation.




Figure 1. $\mu - \delta$ curves obtained with $U/t = 4$. The bold circles lie in the Mott-insulating side of the MIT surface at half filling.

REFERENCES

1. For a review, see M. Imada, A. Fujimori and Y. Tokura, Rev. Mod. Phys. **70** (1998) 1039.
2. A. George *et al.*, Rev. Mod. Phys. **68** (1996) 13.
3. S. Onoda and M. Imada, Phys. Rev. B **67** (2003) R161102, and references therein.
4. S. Onoda and M. Imada, J. Phys. Soc. Jpn. **70** (2001) 632; *ibid.* **70** (2001) 3398.
5. Y.M. Vilk and A.-M.S. Tremblay, J. Phys. (France) I **7** (1997) 1309.
6. H. Matsumoto and F. Mancini, Phys. Rev. B **55** (1997) 2095.
7. F. F. Assaad and M. Imada, Eur. Phys. J. B **10** (1999) 595.
8. C. Castellani *et al.*, Phys. Rev. Lett. **43** (1979) 1957.
9. T. Kashima and M. Imada, J. Phys. Soc. Jpn. **70** (2001) 3052.